



COMPREHENSIVE ANALYSIS OF SILYMARIN
COMPLEX DYNAMICS UNDER SALINITY
STRESS IN *SILYBUM MARIANUM* L.:
INTEGRATIVE BIOCHEMICAL, MOLECULAR,
AND PHYSIOLOGICAL PERSPECTIVES

Olimova Dildora Ikhtiyorovna

Doctoral Researcher, Bukhara State University

Lecturer, Zarmed University

Abstract: Soil salinity is a major environmental constraint that significantly affects plant growth, metabolism, and secondary metabolite biosynthesis. *Silybum marianum* L., an important medicinal plant, produces silymarin—a complex mixture of flavonolignans with potent antioxidant and hepatoprotective properties. The present study provides a comprehensive and integrative analysis of the dynamic behavior of the silymarin complex under varying levels of salt stress. The results demonstrate that salinity induces a dose-dependent modulation of silymarin biosynthesis, characterized by a stimulatory effect at moderate stress levels and inhibitory effects under severe stress conditions. This response is mediated through the activation of the phenylpropanoid pathway, enhanced antioxidant defense mechanisms, and transcriptional regulation of key biosynthetic genes. The study further explores the physiological adaptations, metabolic trade-offs, and stress signaling pathways involved. These findings contribute to a deeper understanding of plant stress physiology and offer practical strategies for enhancing the production of valuable phytochemicals in saline environments.

1. Introduction: Salinity stress is one of the most critical abiotic stress factors limiting agricultural productivity worldwide, affecting more than 800 million hectares of land. The accumulation of soluble salts

in soil leads to osmotic stress, ionic toxicity, nutrient imbalance, and oxidative damage, ultimately impairing plant growth and development. At the cellular level, salt stress disrupts water uptake due to reduced soil water potential, while excessive sodium (Na^+) and chloride (Cl^-) ions interfere with essential metabolic processes. One of the most damaging consequences of salinity is the overproduction of reactive oxygen species (ROS), including superoxide radicals (O_2^-), hydrogen peroxide (H_2O_2), and hydroxyl radicals ($\cdot\text{OH}$), which can damage lipids, proteins, and nucleic acids. Plants have evolved complex adaptive mechanisms to cope with salinity stress, including: osmotic adjustment, ion homeostasis, activation of antioxidant systems, reprogramming of primary and secondary metabolism. Secondary metabolites, particularly phenolic compounds, play a crucial role in plant defense. In this context, *Silybum marianum* L. is of particular interest due to its ability to synthesize silymarin, a biologically active complex of flavonolignans derived from the phenylpropanoid pathway. Silymarin consists mainly of: silybin A and B (dominant isomers), isosilybin, silychristin, silydianin. These compounds exhibit strong antioxidant, anti-inflammatory, and membrane-stabilizing properties, making them valuable in pharmaceutical applications. Despite extensive research on silymarin pharmacology, the regulatory mechanisms governing its biosynthesis under abiotic stress conditions remain insufficiently understood. In particular, the dynamic response of silymarin to salt stress, including its biochemical regulation and physiological significance, requires further investigation.

2. Theoretical Background and Mechanistic Framework. Phenylpropanoid Pathway and Silymarin

Biosynthesis. Silymarin biosynthesis originates from the amino acid phenylalanine through the phenylpropanoid pathway. The key steps include: Phenylalanine → Cinnamic acid (via PAL), Cinnamic acid → p-coumaroyl-CoA, Formation of flavonoid intermediates (via CHS, CHI), Oxidative coupling with coniferyl alcohol → flavonolignans. The regulation of this pathway is tightly controlled at both enzymatic and transcriptional levels.

2.1 Salt Stress Signaling Mechanisms: Salt stress perception triggers multiple signaling pathways: calcium signaling (Ca^{2+} influx), activation of MAP kinases, phytohormone signaling (ABA, jasmonic acid), ROS-mediated signaling cascades. These signals lead to transcriptional activation of stress-responsive genes, including those involved in secondary metabolism.

3. Materials and Methods (Further Detailed).
3.1 Experimental Setup The experiment followed a completely randomized design with four treatments and three biological replicates. Environmental conditions were strictly controlled to minimize variability.

3.2 Sampling Strategy: Samples were collected at three time points: early response (7 days), adaptive phase (14 days), prolonged stress (21 days). This allowed analysis of temporal dynamics.

3.3 Advanced Analytical Techniques: HPLC with UV detection for silymarin profiling Spectrophotometric assays for antioxidant enzymes, qRT-PCR for gene expression Correlation analysis between biochemical parameters

4. Results (Highly Expanded).4.1 Temporal Dynamics of Growth Inhibition

Growth reduction was gradual and time-dependent. Under moderate stress, plants maintained relatively stable growth during early stages, suggesting effective acclimation. However, prolonged exposure led to cumulative damage.

4.2 Time-Dependent Silymarin Accumulation. Silymarin accumulation showed dynamic temporal variation: **Day 7:** slight increase (stress perception phase), **Day 14:** peak accumulation (adaptive phase), **Day 21:** decline under severe stress. This indicates that biosynthesis is tightly linked to stress duration and plant adaptive capacity.

4.3 Metabolic Trade-Offs: Under stress conditions, plants reallocate resources:

- reduced primary metabolism (growth),
- enhanced secondary metabolism (defense).

This trade-off explains increased silymarin accumulation under moderate stress.

4.4 Oxidative Stress Markers

MDA levels increased significantly under severe stress, indicating lipid peroxidation. However, under moderate stress, antioxidant systems effectively controlled ROS levels.

4.5 Enzymatic Antioxidant Network

The coordinated increase in SOD, CAT, and GPX suggests a synergistic defense mechanism. SOD converts superoxide radicals into H_2O_2 , which is then detoxified by CAT and GPX.

5. Discussion (Deep Scientific Interpretation): The findings demonstrate that silymarin biosynthesis is an integral part of the plant's adaptive response to salinity stress. Moderate salt stress enhances metabolic flux through the phenylpropanoid pathway, resulting in increased production of flavonolignans. The role of ROS as signaling molecules is particularly important. While excessive ROS can cause damage, controlled ROS production acts as a trigger for defense responses. This dual role explains the biphasic nature of the response. Furthermore, the decline in silymarin under severe stress reflects a breakdown of metabolic homeostasis. Energy limitations, enzyme inactivation, and damage to cellular structures collectively inhibit biosynthesis.

6. Integrated Model of Silymarin Regulation under Salt Stress: The study proposes the following model: **Salt stress → ROS production → signaling activation → gene expression (PAL, CHS) → increased silymarin → antioxidant protection.** Under excessive stress: **Salt stress → excessive ROS → cellular damage → enzyme inhibition → decreased silymarin.**

Conclusion: This study provides a comprehensive understanding of how salinity affects silymarin biosynthesis in *Silybum marianum*. The results highlight the importance of stress intensity, duration, and metabolic regulation in determining phytochemical accumulation.

References

1. Dixon, R. A., & Paiva, N. L. (1995). Stress-induced phenylpropanoid metabolism. *Plant Cell*.
2. Parida, A. K., & Das, A. B. (2005). Salt tolerance mechanisms. *Ecotoxicology*.

3. ElSayed, A. I., et al. (2019). Stress responses in medicinal plants. *Plants*.
4. Asakereh, F., et al. (2025). Flavonolignan biosynthesis genes. *BMC Plant Biology*.
5. Bahari, H., et al. (2024). Antioxidant role of silymarin. *Inflammopharmacology*.
6. Taiz, L., & Zeiger, E. (Plant Physiology textbook reference).